

SPATIO-TEMPORAL MODELING OF URBAN EXTREME RAINFALL EVENTS AT HIGH RESOLUTION

EGU, Vienna, April 28, 2025

Chloé SERRE-COMBE ¹ Nicolas MEYER ¹ Thomas OPITZ ² Gwladys TOULEMONDE ¹

¹IMAG, Université de Montpellier, LEMON Inria

²INRAE, BioSP, Avignon




GENERAL CONTEXT - MONTPELLIER, FRANCE



Floods in Montpellier, September 2022 and August 2015 (*Midi Libre*)

- Mediterranean events, localized rainfall
- Urban area, flood risks






- **Study area:** Verdanson water catchment
- **Source:** Urban observatory of HydroScience Montpellier (HSM)¹
- **Time period:** [2019, 2023[
- **Temporal resolution:** 5 minutes
- **Spatial resolution:** 77 m to 1531 m

¹FINAUD-GUYOT et al., 2023


$$\mathcal{S} = \{17 \text{ rain gauges}\} \subset \mathbb{R}^2 \text{ and } \mathcal{T} \subset \mathbb{R}_+$$



► Generalized Pareto Distribution (GPD)²

$$X \mid X > u \sim \underbrace{H_\xi}_{\text{GPD}(\xi, \sigma, u)} \text{ with } \xi \in \mathbb{R}, \sigma > 0.$$



²PICKANDS III, 1975




► Extended Generalized Pareto Distribution (EGPD)³

$$X \sim \underbrace{G(H_\xi)}_{\text{EGPD}(\xi, \sigma, \kappa)} \text{ with } G(x) = x^\kappa, \kappa > 0$$

³NAVEAU et al., 2016



EGPD fitting for two rain gauges, CRBM (left) and CNRS (right) with left-censoring and 95% CI



Univariate

Dependence

Rainfall random field: $\mathbf{X} = \{X_{s,t}, (s, t) \in \mathcal{S} \times \mathcal{T}\}$, stationary and isotropic.

Let $\Lambda_{\mathcal{S}} \subset \mathbb{R}_+^2$ and $\Lambda_{\mathcal{T}} \subset \mathbb{R}_+$ be sets of spatial and temporal lags respectively.

Rainfall random field: $\mathbf{X} = \{X_{s,t}, (s, t) \in \mathcal{S} \times \mathcal{T}\}$, stationary and isotropic.

Let $\Lambda_{\mathcal{S}} \subset \mathbb{R}_+^2$ and $\Lambda_{\mathcal{T}} \subset \mathbb{R}_+$ be sets of spatial and temporal lags respectively.

Variogram (MATHERON, 1963)

$$\gamma(\mathbf{h}, \tau) = \frac{1}{2} \text{Var}(X_{s,t} - X_{s+\mathbf{h}, t+\tau})$$

- Quantifies variability
- Higher $\gamma(\mathbf{h}, \tau) \rightarrow$ weaker dependence

$$\mathbf{h} \in \Lambda_{\mathcal{S}}, \tau \in \Lambda_{\mathcal{T}}$$

Rainfall random field: $\mathbf{X} = \{X_{s,t}, (s,t) \in \mathcal{S} \times \mathcal{T}\}$, stationary and isotropic.

Let $\Lambda_{\mathcal{S}} \subset \mathbb{R}_+^2$ and $\Lambda_{\mathcal{T}} \subset \mathbb{R}_+$ be sets of spatial and temporal lags respectively.

Variogram (MATHERON, 1963)

$$\gamma(\mathbf{h}, \tau) = \frac{1}{2} \text{Var}(X_{s,t} - X_{s+\mathbf{h}, t+\tau})$$


- Quantifies variability
- Higher $\gamma(\mathbf{h}, \tau) \rightarrow$ weaker dependence

$\mathbf{h} \in \Lambda_{\mathcal{S}}, \tau \in \Lambda_{\mathcal{T}}, X_{s,t}^*$ uniform margins.

Extremogram (DAVIS and MIKOSCH, 2009)

$$\chi(\mathbf{h}, \tau) = \lim_{q \rightarrow 1} \mathbb{P}(X_{s,t}^* > q \mid X_{s+\mathbf{h}, t+\tau}^* > q)$$

- Measures tail dependence
- Higher $\chi(\mathbf{h}, \tau) \rightarrow$ stronger dependence




Definition (de FONDEVILLE and DAVISON, 2018)

For all $s \in \mathcal{S}$ and $t \in \mathcal{T}$, a risk function $r(\mathbf{X}) = X_{s_0, t_0}$,

$$u^{-1} X_{s,t} \mid X_{s_0, t_0} > u \xrightarrow{d} Y_{s,t} \quad \text{with} \quad Y_{s,t} = R_{s,t} e^{W_{s,t} - W_{s_0, t_0} - \gamma(s - s_0, t - t_0)},$$

where (s_0, t_0) is a space-time location, u is a high threshold, $R_{s,t} \sim \text{Pareto}(1)$, $W_{s,t}$ is a Gaussian process.



Dependence measures

► r -variogram:

$$\gamma_r(\mathbf{h}, \tau) = \frac{1}{2} \text{Var}(W_{s_0, t_0} - W_{s_0 + \mathbf{h}, t_0 + \tau})$$

► r -extremogram:

$$\chi_r(\mathbf{h}, \tau) = \lim_{q \rightarrow 1} \mathbb{P}(X_{s_0 + \mathbf{h}, t_0 + \tau}^* > q)$$

Spatio-temporal extremogram with a Brown-Resnick dependence

Let $\mathbf{h} \in \Lambda_S$ and $\tau \in \Lambda_T$. We have

$$\chi(\mathbf{h}, \tau) = 2 \left(1 - \phi \left(\sqrt{\frac{1}{2} \gamma(\mathbf{h}, \tau)} \right) \right)$$

with ϕ the std normal c.d.f. and γ the variogram of $\mathbf{W}$.

Spatio-temporal extremogram with a Brown-Resnick dependence

Let $\mathbf{h} \in \Lambda_S$ and $\tau \in \Lambda_T$. We have

$$\chi(\mathbf{h}, \tau) = 2 \left(1 - \phi \left(\sqrt{\frac{1}{2} \gamma(\mathbf{h}, \tau)} \right) \right)$$

with ϕ the std normal c.d.f. and γ the variogram of $\mathbf{W}$.

Separable model: Fractional Brownian motion with additive separability.

$$\frac{\gamma(\mathbf{h}, \tau)}{2} = \beta_1 \|\mathbf{h}\|^{\alpha_1} + \beta_2 |\tau|^{\alpha_2}$$

Spatio-temporal extremogram with a Brown-Resnick dependence

Let $\mathbf{h} \in \Lambda_S$ and $\tau \in \Lambda_T$. We have

$$\chi(\mathbf{h}, \tau) = 2 \left(1 - \phi \left(\sqrt{\frac{1}{2} \gamma(\mathbf{h}, \tau)} \right) \right)$$

with ϕ the std normal c.d.f. and γ the variogram of $\mathbf{W}$.

Separable model: Fractional Brownian motion with additive separability.

$$\frac{\gamma(\mathbf{h}, \tau)}{2} = \beta_1 \|\mathbf{h}\|^{\alpha_1} + \beta_2 |\tau|^{\alpha_2}$$

- Efficient estimation using WLSE (BUHL et al., 2019)

Spatio-temporal extremogram with a Brown-Resnick dependence

Let $\mathbf{h} \in \Lambda_S$ and $\tau \in \Lambda_T$. We have

$$\chi(\mathbf{h}, \tau) = 2 \left(1 - \phi \left(\sqrt{\frac{1}{2} \gamma(\mathbf{h}, \tau)} \right) \right)$$


with ϕ the std normal c.d.f. and γ the variogram of $\mathbf{W}$.

Separable model: Fractional Brownian motion with additive separability.


$$\frac{\gamma(\mathbf{h}, \tau)}{2} = \beta_1 \|\mathbf{h}\|^{\alpha_1} + \beta_2 |\tau|^{\alpha_2}$$

- Efficient estimation using WLSE (BUHL et al., 2019)
- **Not realistic**


Separable model




Separable model




Separable model



Separable model



Non-separable model




Towards more realistic modeling: introduce advection $\mathbf{V}$ to relax separability

$$\gamma_L(\mathbf{h}, \tau) = \gamma(\mathbf{h} - \tau \mathbf{V}, \tau)$$

$$\Rightarrow \frac{1}{2} \gamma_L(\mathbf{h}, \tau) = \beta_1 \|\mathbf{h} - \tau \mathbf{V}\|^{\alpha_1} + \beta_2 |\tau|^{\alpha_2}$$

► Parameters: $\Theta = (\beta_1, \beta_2, \alpha_1, \alpha_2, \mathbf{V})$

Non-separable model



Goal: Estimate variogram parameters $\Theta = (\beta_1, \beta_2, \alpha_1, \alpha_2, \mathbf{V})$

Joint exceedances count

$$K_{\mathbf{h}, \tau}(u) = \sum_{(\mathbf{s}_i, \mathbf{s}_j) \in \mathcal{N}(\mathbf{h})} \sum_{(t_i, t_j) \in \mathcal{N}(\tau)} \mathbb{1}_{\{X_{\mathbf{s}_i, t_i} > u, X_{\mathbf{s}_j, t_j} > u\}}$$

Asymptotic distribution (large u) on r -Pareto:

$$K_{\mathbf{h}, \tau} \sim \mathcal{B} \left(\sum_{(\mathbf{s}_0, t_0) \in \mathcal{R}} \#\{\mathbf{s} : (\mathbf{s}, \mathbf{s}_0) \in \mathcal{N}(\mathbf{h})\}, \chi_{r, \Theta}(\mathbf{h}, \tau) \right)$$

with

- ▶ $\mathcal{R}$ is a subset of spatio-temporal conditioning points
- ▶ $\mathcal{N}(\mathbf{h})$ the set of pairs of sites separated by $\mathbf{h}$
- ▶ $\mathcal{N}(\tau)$ the set of pairs of times separated by τ

Asymptotic distribution (large u) on r -Pareto:

$$K_{\mathbf{h},\tau} \sim \mathcal{B} \left(\sum_{(\mathbf{s}_0, t_0) \in \mathcal{R}} \#\{\mathbf{s} : (\mathbf{s}, \mathbf{s}_0) \in \mathcal{N}(\mathbf{h})\}, \chi_{r,\Theta}(\mathbf{h}, \tau) \right)$$


Composite log-likelihood maximization

$$\ell_C(\Theta) \propto \sum_{(\mathbf{s}_0, t_0) \in \mathcal{R}} \sum_{(\mathbf{s}, \mathbf{s}_0) \in \mathcal{N}(\mathbf{h})} \sum_{(t, t_0) \in \mathcal{N}(\tau)} k_{\mathbf{s},t} \log \chi_{r,\Theta}(\mathbf{h}, \tau) + (1 - k_{\mathbf{s},t}) \log(1 - \chi_{r,\Theta}(\mathbf{h}, \tau))$$


with $k_{\mathbf{s},t} = \mathbb{1}_{\{X_{\mathbf{s},t} > u, X_{\mathbf{s}_0, t_0} > u\}}$, for large u

- Optimization: maximization of $\ell_C(\Theta)$ with respect to Θ

VALIDATION ON SIMULATIONS





Without advection




With advection



Parameter estimation on 100 simulations of 1000 replicates of r -Pareto processes with 25 sites and 30 time observations








- BUHL, S., DAVIS, R. A., KLÜPPELBERG, C., & STEINKOHL, C. (2019). Semiparametric estimation for isotropic max-stable space-time processes.
- DAVIS, R. A., & MIKOSCH, T. (2009). The extremogram: A correlogram for extreme events.
- de FONDEVILLE, R., & DAVISON, A. C. (2018). High-dimensional peaks-over-threshold inference. *Biometrika*, 105(3), 575–592.
- FINAUD-GUYOT, P., GUINOT, V., MARCHAND, P., NEPPEL, L., SALLES, C., & TOULEMONDE, G. (2023). Rainfall data collected by the HSM urban observatory (OMSEV).
- MATHERON, G. (1963). Principles of geostatistics. *Economic geology*, 58(8), 1246–1266.
- NAVEAU, P., HUSER, R., RIBEREAU, P., & HANNART, A. (2016). Modeling jointly low, moderate, and heavy rainfall intensities without a threshold selection. *Water Resources Research*.
- PICKANDS III, J. (1975). Statistical inference using extreme order statistics. *the Annals of Statistics*, 119–131.



Rainfall amounts on CNRS and Polytech rain gauges

Pixels of 1km x 1km



- **Source:** COMEPHORE, Météo France
- **Time period:** [1997, 2023[
- **Temporal resolution:** Every hour
- **Spatial resolution:** 1 km²

More consistent data: Both datasets + Neural Network Downscaling.

Generalized Pareto Distribution



Extended GPD⁴

$$\overline{H}_\xi \left(\frac{x-u}{\sigma} \right) = \begin{cases} (1 + \xi \frac{x-u}{\sigma})_+^{-1/\xi} & \text{if } \xi \neq 0, \\ e^{-\frac{x-u}{\sigma}} & \text{if } \xi = 0, \end{cases}$$

where $a_+ = \max(a, 0)$, $\sigma > 0$, $x - u > 0$

- Models extreme precipitation
- Depends on a threshold choice


$$F(x) = G \left(H_\xi \left(\frac{x}{\sigma} \right) \right),$$

where $G(x) = x^\kappa$, $\kappa > 0$

- Models **moderate and extreme** precipitation
- Avoids a threshold choice


⁴NAVEAU et al., 2016

Case of additive separability: $\frac{\gamma(h, \tau)}{2} = \beta_1 \|h\|^{\alpha_1} + \beta_2 |\tau|^{\alpha_2}, \quad 0 < \alpha_1, \alpha_2 \leq 2, \quad \beta_1, \beta_2 > 0$



ESTIMATION OF THE SEPARABLE VARIOGRAM PARAMETERS

Case of additive separability: $\frac{\gamma(h, \tau)}{2} = \beta_1 \|h\|^{\alpha_1} + \beta_2 |\tau|^{\alpha_2}, \quad 0 < \alpha_1, \alpha_2 \leq 2, \quad \beta_1, \beta_2 > 0$



Weighted Least Squares Estimation (WLSE)

$$\begin{pmatrix} \hat{c}_i \\ \hat{\alpha}_i \end{pmatrix} = \operatorname{argmin}_{c_i, \alpha_i} \sum_x w_x \left(\eta(\hat{\chi}) - (c_i + \alpha_i x) \right)^2$$


Empirical spatial extremogram

For a fixed $t \in \mathcal{T}$ and q a high quantile,

$$\hat{\chi}_q^{(t)}(\mathbf{h}, 0) = \frac{\frac{1}{|N_{\mathbf{h}}|} \sum_{i,j \mid (s_i, s_j) \in N_{\mathbf{h}}} \mathbb{1}_{\{X_{s_i, t}^* > q, X_{s_j, t}^* > q\}}}{\frac{1}{|S|} \sum_{i=1}^{|S|} \mathbb{1}_{\{X_{s_i, t}^* > q\}}},$$


where $C_{\mathbf{h}}$ are equifrequent distance classes and

$$N_{\mathbf{h}} = \left\{ (s_i, s_j) \in S^2 \mid \|s_i - s_j\| \in C_{\mathbf{h}} \right\}.$$



Transformation and WLSE

$\Rightarrow$




Spatial variogram $\hat{\gamma}(\mathbf{h}, 0) = 2\hat{\beta}_1 \|\mathbf{h}\|^{\hat{\alpha}_1}$


Empirical temporal extremogram

For a location $s \in S$, a high quantile q and $t_k \in \{t_1, \dots, t_T\}$,

$$\hat{\chi}_q^{(s)}(\mathbf{0}, \tau) = \frac{\frac{1}{T-\tau} \sum_{k=1}^{T-\tau} \mathbb{1}_{\{X_{s,t_k}^* > q, X_{s,t_k+\tau}^* > q\}}}{\frac{1}{T} \sum_{k=1}^T \mathbb{1}_{\{X_{s,t_k}^* > q\}}}$$



Transformation and WLSE




Temporal variogram $\hat{\gamma}(\mathbf{0}, \tau) = 2\hat{\beta}_2|\tau|^{\hat{\alpha}_2}$

Composite log-likelihood maximization

$$\ell_C(\Theta) \propto \sum_{(s_0, t_0) \in \mathcal{R}} \sum_{(s, s_0) \in \mathcal{N}(h)} \sum_{(t, t_0) \in \mathcal{N}(\tau)} k_{s,t} \log \chi_{r, \Theta}(h, \tau) + (1 - k_{s,t}) \log(1 - \chi_{r, \Theta}(h, \tau))$$

with $k_{s,t} = \mathbb{1}_{\{X_{s,t} > u, X_{s_0, t_0} > u\}}$, for large u



Wind in t_0

Wind-informed advection:

$$\mathbf{V} = \eta_1 \cdot |\mathbf{V}_w|^{\eta_2} \cdot \text{sign}(\mathbf{V}_w), \quad \eta_1, \eta_2 > 0$$

$$\Rightarrow \Theta = (\beta_1, \beta_2, \alpha_1, \alpha_2, \eta_1, \eta_2)$$

with $\mathbf{V}_w$ real wind data.